Archimedes Principle


F=mg F=pVg

ARCHIMEDES' PRINCIPLE

If any object is submerged or partially submerged in a liquid, the body is buoyed up by a force equal to the weight of the liquid displaced by the object.

1)

A 200-kg concrete block has a volume of 0.10 m³. (a) Find the mass of water displaced by this block ($\rho_{\text{water}} = 1000 \text{ kg/m}^3$). (b) Calculate the buoyancy force of the water. (c) Find the force needed to lift the block while it is submerged in water.




2000 kg/ms

a) 2 2000 = 2000 kg

2)

A ship model floats in water as shown. The total volume of the model is 0.25 m^3 , and $6\overline{0}$ percent of this volume is submerged. (a) Find the mass of the water displaced. (b) Calculate the buoyancy force of the water. (c) Find the weight and mass of the ship.



a) $p \cdot V = m$ m = 1000 (60% of 0.25)= 6000 (.15)

= **15**0 kg

b) 18= mg =150(10) =1500N

C)

3)


The density of gasoline is 680 kilograms per cubic meter. A 48-kg ball whose volume is 0.085 m⁸ is tethered to the bottom of a tank of gasoline, as shown. (a) Find the mass of gasoline displaced by this ball, (b) Calculate the buoyancy force of the gasoline, (c) Find the tension in the rope.



アタト デーデョウタト でラトで 手歩ーデョメ

Archimedes Principle

A submerged stone weighs 248 N, but out of the water at weighs 688 N. (a) Calculate the buoyancy force of the water. (b) I ind the mass of the water displaced, ic) Use this mass and p_{water} = 1000 kg/m² to find the volume of the displaced water, which will tell you the volume of the stone.



5)

A 2000-kg homogeneous beam supports a mass of 10.0 kg, as shown Find the tension in the wire. Then find the vertical and horizontal forces at the wall.

